CANADIAN JOURNAL OF FOREST RESEARCH

JOURNAL CANADIEN DE RECHERCHE FORESTIÈRE

Volume 17, 1987

Author Index / Index des auteurs

Ackerley, C.A. see Massicotte, H.B., 846
Adams, W.T. see Merkle, S.A., 402
Agee, J.K., and Huff, M.H. Fuel succession in a western hemlock/Douglas-fir forest, 697
Alden, J., and Loopstra, C. Genetic diversity and population structure of *Picea glauca* on an altitudinal gradient in interior Alaska, 1519
Alli, I. see Jones, A.R.C., 263
Amaranthus, M.P. see Perry, D.A., 929
Amarantus, M.P., and Perry, D.A. Effect of soil transfer on ectomycorrhiza formation and the survival and growth of conifer seedlings on old, nonreforested clear-cuts, 944
Amman, G.D. see McGregor, M.D., 1234
Andersen, C.P., Sucoff, E.I., and Dixon, R.K. The influence of low soil temperature on the growth of vesicular–arbuscular mycorrhizal *Fraxinus pennsylvanica*, 951
Andersson, B. see Nilsson, J.-E., 1340
Anttila, P. see Raunemaa, T., 466
Armstrong, G. see Timmer, V.R., 644
Ashbee, D. see Hunt, R.S., 594
Atalay, A. see Mitchell, R.J., 1153
Atkinson, C.J. see Denne, M.P., 1166
Auchmoody, L.R. Evaluating growth responses to fertilization: Reply, 94
Bailey, R.L. see McTague, J.P., 87
Barclay, H.J. see Pang, P.C., 1379
Barclay, J.L. see Tippett, J.T., 1228
Bare, B.B. see Rustagi, K.P., 1401
Barnes, B.V. see Denton, S.R., 598
Barnes, B.V. see Denton, S.R., 613
Barrett, J.W., Knowles, P., and Cheliak, W.M. The mating system in a black spruce clonal seed orchard, 379
Barrett, J.W., Cheliak, W.M., and Knowles, P.K. Variation in the PGI/AAT linkage group between populations of black spruce, 756
Barrière, G. see Comps, B., 1043
Bartsch, N. Responses of root systems of young *Pinus sylvestris* and *Picea abies* plants to water deficits and soil acidity, 805
Baser, C.M., Garrett, H.E., Mitchell, R.J., Cox, G.S., and Starbuck, C.J. Indolebutyric acid and ectomycorrhizal inoculation increase lateral root initiation and development of container-grown black oak seedings, 36
Bauce, E. see Gagnon, R.R., 105
Bauer, E.O. see Cecich, R.A., 1408
Beaulieu, J. see Corriveau, A., 675
Becker, C.A., Mroz, G.D., and Fuller, L.G. The effects of plant moisture stress on red pine (*Pinus resinosa*) seedling growth and establishment, 813
Bedker, P.J., Wingfield, M.J., and Blanchette, R.A. Pathogenicity of *Bursaphelenchus xylophilus* on three species of pine, 51
Bella, I.E., and Navratil, S. Growth losses from winter drying (red belt damage) in lodgepole pine stands on the east slopes of the Rockies in Alberta, 1289
Benson, M.L. see Linder, S., 1157
Bergeron, Y., and Gagnon, D. Age structure of red pine (*Pinus resinosa* Ait.) at its northern limit in Quebec, 129
Bernier, P.Y. Regeneration and growth of lodgepole pine in small openings in the Alberta Foothills, 758
Bertrand, F. see Ung, C.-H., 588
Bidlake, W.R. see Higgins, S.S., 1364
Black, R.A. see Higgins, S.S., 1364
Black, T.A. see Livingston, N.J., 1115
Black, T.A. see Livingston, N.J., 1273
Blanche, C.A. see Paine, T.D., 1202
Blanchette, R.A. see Bedker, P.J., 51
Bonga, J.M. see von Aderkas, P., 1293
Boninsegna, J.A. see Villalba, R., 1527
Borden, J.H., Ryker, L.C., Chong, L.J., Pierce, H.D., Jr., Johnston, B.D., and Oehlschlager, A.C. Response of the mountain pine beetle, *Dendroctonus ponderosae* Hopkins (Coleoptera: Scolytidae), to five semiochemicals in British Columbia lodgepole pine forests, 118
Borden, J.H. see Brooks, J.E., 740
Borden, J.H. see Maclauchlan, L.E., 1124
Bousquet, J., Cheliak, W.M., and Lalonde, M. Genetic differentiation among 22 mature populations of green alder (*Alnus crispa*) in central Quebec, 219

Bowen, G.D. see Garbaye, J., 941
Boyle, C.D., Robertson, W.J., and Salonius, P.O. Use of mycelial slurries of mycorrhizal fungi as inoculum for commercial tree seedling nurseries, 1480
Bräker, O.U. see Kienast, F., 683
Brand, D.G. Estimating the surface area of spruce and pine foliage from displaced volume and length, 1305
Bridgwater, F.E. see Brown, S.D., 299
Brisbin, R.L., and Dale, M.E. Estimating tree-quality potential in a managed white oak stand by Markov Chain analysis, 9
Brooks, J.E., Borden, J.H., and Pierce, H.D., Jr. Foliar and cortical monoterpenes in Stika spruce: potential indicators of resistance to the white pine weevil, *Pissodes strobi* Peck (Coleoptera: Curculionidae), 740
Brown, J.K., and DeByle, N.V. Fire damage, mortality, and suckering in aspen, 1100
Brown, S.D., and Bridgwater, F.E. Observations on pollination in loblolly pine, 299
Browne, B. see Hughes, J.W., 1623
Burdett, A.N. Understanding root growth capacity: theoretical considerations in assessing planting stock quality by means of root growth tests, 768
Burgess, D.M. see Hendrickson, O.Q., 210
Burkhart, H.E. see Sprinz, P.T., 534
Burkhart, H.E. see Dyer, M.E., 572
Byrne, S.V., Wentworth, T.R., and Nusser, S.M. A moisture strain index for loblolly pine, 23
Cackette, M.R. see Maclauchlan, L.E., 1124
Camiré, C. see Côté, B., 515
Cao, Q.V. see Parresol, B.R., 250
Carlson, W.C., and Harrington, C.A. Cross-sectional relationships in root systems of loblolly and shortleaf pine, 556
Castellano, M.A. see Entry, J.A., 865
Cecich, R.A., and Bauer, E.O. The acceleration of jack pine seed development, 1408
Ceulemans, R., Impens, I., and Steenackers, V. Variations in photosynthetic, anatomical, and enzymatic leaf traits and correlations with growth in recently selected *Populus* hybrids, 273
Chatarpaul, L. see Hendrickson, O.Q., 210
Cheliak, W.M. see Bousquet, J., 219
Cheliak, W.M. see Barrett, J.W., 379
Cheliak, W.M. see Barrett, J.W., 756
Chong, L.J. see Borden, J.H., 118
Choquette, C. see Perry, D.A., 1434
Cline, S.P. see Sollins, P., 1585
Clyde, M.A., and Titus, S.J. Radial and longitudinal variation in stem diameter increment of lodgepole pine, white spruce, and black spruce: species and crown class differences, 1223
Cole, E.C., and Newton, M. Fifth-year responses of Douglas-fir to crowding and nonconiferous competition, 181
Colombo, S.J. Changes in osmotic potential, cell elasticity, and turgor relationships of 2nd-year black spruce container seedlings, 365
Colombo, S.J. Erratum: Changes in osmotic potential, cell elasticity, and turgor relationships of 2nd-year black spruce container seedlings, 1312
Colombo, S.J. see Kim, Y.T., 27
Comps, B., Barrière, G., Merzeau, D., et Letouzey, J. La variabilité alloenzymatique des hêtraies dans les sous-domaines medio- et eu-atlantique d'Europe, 1043
Connor, K.F., and Lanner, R.M. The architectural significance of interfoliar branches in *Pinus* subsection *Balfourianae*, 269
Corriveau, A., Beaulieu, J., and Mothe, F. Wood density of natural white spruce populations in Quebec, 675
Côté, B., and Camiré, C. Tree growth and nutrient cycling in dense plantings of hybrid poplar and black alder, 515
Coutts, M.P. Developmental processes in tree root systems, 761
Cox, G.S. see Baser, C.M., 36
Cox, G.S. see Mitchell, R.J., 1153
Critchfield, W.B. Allozyme variation in *Picea mariana* from Newfoundland: Discussion, 1471
Cromack, K., Jr. see Entry, J.A., 865
Cromack, K., Jr. see Harmon, M.E., 1265
Cromack, K., Jr. see Entry, J.A., 260
Cropper, W.P., Jr. see Ewel, K.C., 325
Cropper, W.P., Jr. see Ewel, K.C., 330
Cunia, T. On the error of continuous forest inventory estimates, 436
Cunia, T. On the error of estimates from successive forest inventories with SPR, 442
Dale, M.E. see Brisbin, R.L., 9
Dale, V.H. see Pastor, J., 1394
Dancik, B.P. see Williams, D.J., 1442
D'Auria, J.M. see Maclauchlan, L.E., 1124
David, M.B., and Gertner, G.Z. Sources of variation in soil solution collected by tension plate lysimeters, 190
Davis, D.D. see Umbach, D.M., 1213
DeByle, N.V. see Brown, J.K., 1100

DeLucia, E.H., and Smith, W.K. Air and soil temperature limitations on photosynthesis in Engelmann spruce during summer, 527
Denne, M.P., and Atkinson, C.J. Reactivation of vessel expansion in relation to budbreak in sycamore (*Acer pseudoplatanus*) trees, 1166
Denton, S.R., and Barnes, B.V. Spatial distribution of ecologically applicable climatic statistics in Michigan, 598
Denton, S.R., and Barnes, B.V. Tree species distributions related to climatic patterns in Michigan, 613
Desponts, M., et Simon, J.-P. Structure et variabilité génétique de populations d'épinette noire (*Picea mariana* (Mill.) B.S.P.) dans la zone hémiarctique du Nouveau -Québec, 1006
Desprez-Loustau, M.L., et Dessureault, M. Étiologie du chancre godronien du bouleau jaune, 1355
Dessureault, M. see Desprez-Loustau, M.L., 1355
DeWald, L.E., and Feret, P.P. Changes in loblolly pine root growth potential from September to April, 635
Dixon, R.K. see Andersen, C.P., 951
Dixon, R.K. see Mitchell, R.J., 1153
Donahue, J.P., and Howard, A.F. Hydraulic design of culverts on forest roads, 1545
Doucet, R., and Veilleux, J.-M. Evaluating growth response to fertilization: Discussion, 93
Dupper, G.E. see Kinloch, B.B., Jr., 448
Duryea, M.L., and Omi, S.K. Top pruning Douglas-fir seedlings: morphology, physiology, and field performance, 1371
Durzan, D.J. see Gupta, P.K., 1130
Dyer, M.E., and Burkhart, H.E. Compatible crown ratio and crown height models, 572
Eberhart, K.E., and Woodard, P.M. Distribution of residual vegetation associated with large fires in Alberta, 1207
Edmonds, R.L. Decomposition rates and nutrient dynamics in small-diameter woody litter in four forest ecosystems in Washington, U.S.A., 499
Eis, S. Root systems of older immature hemlock, cedar, and Douglas-fir, 1348
Ek, A.R. see Payandeh, B., 95
Entry, J.A., and Cromack, K., Jr. Effect of pH, aluminum, and sulfate on a high-elevation *Armillaria* isolate cultured in vitro, 260
Entry, J.A., Stark, N.M., and Loewenstein, H. Effect of timber harvesting on extractable nutrients in a northern Rocky Mountain forest soil, 735
Entry, J.A., Cromack, K., Jr., Stafford, S.G., and Castellano, M.A. The effect of pH and aluminum concentration on ectomycorrhizal formation in *Abies balsamea*, 865
Errico, D. see Reed, W.J., 1455
Espinosa Bancalari, M.A., Perry, D.A., and Marshall, J.D. Leaf area – sapwood area relationships in adjacent young Douglas-fir stands with different early growth rates, 174
Espinosa Bancalari, M.A., and Perry, D.A. Distribution and increment of biomass in adjacent young Douglas-fir stands with different early growth rates, 722
Ewel, K.C., Cropper, W.P., Jr., and Gholz, H.L. Soil CO_2 evolution in Florida slash pine plantations. I. Changes through time, 325
Ewel, K.C., Cropper, W.P., Jr., and Gholz, H.L. Soil CO_2 evolution in Florida slash pine plantations. II. Importance of root respiration, 330
Fahey, T.J. see Hughes, J.W., 1623
Federer, C.A., and Hornbeck, J.W. Expected decrease in diameter growth of even-aged red spruce, 266
Feret, P.P. see DeWald, L.E., 635
Ferm, A. see Rinne, P., 545
Finkle, B.J. see Gupta, P.K., 1130
Fisher, J.T., and Neumann, R.W. Cultivation and weed control for aspen seedling establishment in the southern Rocky Mountains, 47
Fortin, J.A. see Gagnon, J., 840
Fortin, J.A. see McAfee, B.J., 859
Foster, G.S., Lambeth, C.C., and Greenwood, M.S. Growth of loblolly pine rooted cuttings compared with seedlings, 157
Fowler, D.P. The hybrid white × Sitka spruce: species crossability, 413
Fowler, D.P. In defense of the polycross, 1624
Fowler, D.P. see Park, Y.S., 1175
Frandsen, W.H. The influence of moisture and mineral soil on the combustion limits of smoldering forest duff, 1540
Fuller, L.G. see Becker, C.A., 813
Fyles, J.W., and McGill, W.B. Decomposition of boreal forest litters from central Alberta under laboratory conditions, 109
Fyles, J.W., and McGill, W.B. Nitrogen mineralization in forest soil profiles from central Alberta, 242
Gagnon, D. see Bergeron, Y., 129
Gagnon, J., Langlois, C.G., and Fortin, J.A. Growth of containerized jack pine seedlings inoculated with different ectomycorrhizal fungi under a controlled fertilization schedule, 840
Gagnon, R.R., Bauce, E., and Pineau, M. Relation between air water potential and cambial electrical resistance of balsam fir and white spruce after budbreak, 105
Gale, M.R., and Grigal, D.F. Vertical root distributions of northern tree species in relation to succesional status, 829
Garbaye, J., and Bowen, G.D. Effect of different microflora on the success of ectomycorrhizal inoculation of *Pinus radiata*, 941
Gardner, R.H. see Pastor, J., 1394
Garrett, H.E. see Baser, C.M., 36
Garrett, H.E. see Mitchell, R.J., 1153
Genys, J.B. Provenance variation among different populations of *Pinus strobus* from Canada and the United States, 228
Gertner, G.Z. see David, M.B., 190
Gholz, H.L. see Ewel, K.C., 325
Gholz, H.L. see Ewel, K.C., 330

Gilbert, D.E., Johnson, B.R., and Zala, C. A reliability study of the lightning locating network in British Columbia, 1060
Gillingham, M.P., Speyer, M.R., Northway, S., and McLaughlin, R. Feeding preference and its relation to herbivore repellent studies, 146
Glerum, C. see Kim, Y.T., 27
Goodell, B., Hosli, J.P., and Kropp, B. The diffusion and toxicity of the fumigant chloripicrin injected in sugar maple and white birch trees, 1552
Gordon, A.M., Schlentner, R.E., and Van Cleve, K. Seasonal patterns of soil respiration and CO_2 evolution following harvesting in the white spruce forests of interior Alaska, 304
Gower, K.T., Grier, C.C., Vogt, D.J., and Vogt, K.A. Allometric relations of deciduous (*Larix occidentalis*) and evergreen conifers (*Pinus contorta* and *Pseudotsuga menziesii*) of the Cascade Mountains in central Washington, 630
Grace, J.C. see Santantonio, D., 900
Graham, R.T. see Harvey, A.E., 58
Graham, R.T. see Jurgensen, M.F., 1283
Greenwood, M.S. see Foster, G.S., 157
Gref, R., and Tenow, O. Resin acid variation in sun and shade needles of Scots pine (*Pinus sylvestris* L.), 346
Grier, C.C. see Gower, K.T., 630
Grigal, D.F. see Strong, T.F., 705
Grigal, D.F. see Gale, M.R., 829
Grossnickle, S.C. Influence of flooding and soil temperature on the water relations and morphological development of cold-stored black spruce and white spruce seedlings, 821
Gunn, E.A., and Rai, A.K. Modelling and decomposition for planning long-term forest harvesting in an integrated industry structure, 1507
Gupta, P.K., Durzan, D.J., and Finkle, B.J. Somatic polyembryogenesis in embryogenic cell masses of *Picea abies* (Norway spruce) and *Pinus taeda* (loblolly pine) after thawing from liquid nitrogen, 1130
Gupta, P.K. see Khuspe, S.S., 1361
Hällgren, J.-E. see Lundmark, T., 1197
Hamlin, D., and Leary, R. Methods for using an integro-differential equation as a model of tree height growth, 353
Hanley, T.A. Physical and chemical response of understory vegetation to deer use in southeastern Alaska, 195
Hann, D.W., Walters, D.K., and Scrivani, J.A. Incorporating crown ratio into prediction equations for Douglas-fir stem volume, 17
Hann, D.W. see Maguire, D.A., 236
Hansen, G.D., and Nyland, R.D. Effects of diameter distribution on the growth of simulated unevenaged sugar maple stands, 1
Hanus, D. see Rohr, R., 418
Hard, J.S. Vulnerability of white spruce with slowly expanding lower boles on dry, cold sites to early seasonal attack by spruce beetles in south central Alaska, 428
Hari, P. see Raunemaa, T., 466
Harmon, M.E. The influence of litter and humus accumulations and canopy openness on *Picea sitchensis* (Bong.) Carr. and *Tsuga heterophylla* (Raf.) Sarg. seedlings growing on logs, 1475
Harmon, M.E., Cromack, K., Jr., and Smith, B.G. Coarse woody debris in mixed-conifer forests, Sequoia National Park, California, 1265
Harou, P. On a social discount rate for forestry: Reply, 1473
Harrington, C.A. see Carlson, W.C., 556
Harrington, J.B. Climatic change: a review of causes, 1313
Harris, A.S. see Shaw, C.G., III, 334
Harry, D.E. Shoot elongation and growth plasticity in incense-cedar, 484
Harvey, A.E., Jurgensen, M.F., Larsen, M.J., and Graham, R.T. Relationships among soil microsite, ectomycorrhizae, and natural conifer regeneration of old-growth forests in Western Montana, 58
Harvey, A.E. see Jurgensen, M.F., 1283
Hazenberg, G. see Yang, K.C., 648
Hendrickson, O.Q., Burgess, D.M., and Chatarpaul, L. Biomass and nutrients in Great Lakes – St. Lawrence forest species: implications for whole-tree and conventional harvest, 210
Higgins, S.S., Black, R.A., Radamaker, G.K., and Bidlake, W.R. Gas exchange characteristics and water relations of *Larix occidentalis*, 1364
Hinesley, L.E. see Warren, S.L., 1092
Ho, I. Comparison of eight *Pisolithus tinctorius* isolates for growth rate, enzyme activity, and phytohormone production, 31
Ho, I. Enzyme activity and phytohormone production of a mycorrhizal fungus, *Laccaria laccata*, 855
Hobbs, S.D., Stafford, S.G., and Slagle, R.L. Undercutting conifer seedlings: effect on morphology and field performance on droughty sites, 40
Hoff, R.J. Dormancy in *Pinus monticola* seed related to stratification time, seed coat, and genetics, 294
Högberg, K.-A. Needle conductance and root egress of containerized Scots pine and Norway spruce seedlings after surface and deep planting, 783
Hook, D.D. see McKevlin, M.R., 1257
Hook, D.D. see McKevlin, M.R., 1572
Hope, S.M. Classification of decayed *Abies amabilis* logs, 559
Hornbeck, J.W. see Federer, C.A., 266
Hosli, J.P. see Goodell, B., 1552
Hotvedt, J.E. see Parresol, B.R., 250
Howard, A.F. see Donahue, J.P., 1545
Huff, M.H. see Agee, J.K., 697

Hughes, J.W., Fahey, T.J., and Browne, B. A better seed and litter trap, 1623
Hughes, T.F. see Tappeiner, J.C., II, 1300
Hungerford, R.D. Estimation of foliage area in dense Montana lodgepole pine stands, 320
Hungerford, R.D. see Running, S.W., 472
Hunt, R.S., Ying, C.C., and Ashbee, D. Variation in damage among *Pinus contorta* provenances caused by the needle cast fungus *Lophodermella concolor*, 594
Hunt, R.S. see Ying, C.C., 1596
Hyun, J.O., Rajora, O.P., and Zsuffa, L. Genetic variation in trembling aspen in Ontario based on isozyme studies, 1134
Impens, I. see Ceulemans, R., 273
Ingleby, K. see Wilson, J., 957
Jahraus, K. see Marshall, P.L., 1240
James, D.F. see Smith, V.G., 1080
Johnson, B.R. see Gilbert, D.E., 1060
Johnston, B.D. see Borden, J.H., 118
Jones, A.R.C., and Alli, I. Sap yields, sugar content, and soluble carbohydrates of saps and syrups of some Canadian birch and maple species, 263
Jones, R.H., and Raynal, D.J. Root sprouting in American beech: production, survival, and the effect of parent tree vigor, 539
Joyce, D.G. Phenotypic variation in predetermined and free growth in sapling size western larch, 200
Jozsa, L.A., and Powell, J.M. Some climatic aspects of biomass productivity of white spruce stem wood, 1075
Jurgensen, M.F., Larsen, M.J., Graham, R.T., and Harvey, A.E. Nitrogen fixation in woody residue of northern Rocky Mountain conifer forests, 1283
Jurgensen, M.F. see Harvey, A.E., 58
Kabzems, R.D., and Klinka, K. Initial quantitative characterization of soil nutrient regimes. I. Soil properties, 1557
Kabzems, R.D., and Klinka, K. Initial quantitative characterization of soil nutrient regimes. II. Relationships among soils, vegetation, and site index, 1565
Katainen, H.-S. see Raunemaa, T., 466
Kauppi, A. see Rinne, P., 545
Keane, M.G., and Weetman, G.F. Leaf area – sapwood cross-sectional area relationships in repressed stands of lodgepole pine, 205
Khuspe, S.S., Gupta, P.K., Kulkarni, D.K., Mehta, U., and Mascarenhas, A.F. Increased biomass production by tissue culture of eucalyptus, 1361
Kienast, F., Schweingruber, F.H., Bräker, O.U., and Schär, E. Tree-ring studies on conifers along ecological gradients and the potential of single-year analyses, 683
Kim, Y.T., Glerum, C., Stoddart, J., and Colombo, S.J. Effect of fertilization of free amino acid concentrations in black spruce and jack pine containerized seedlings, 27
Kimmins, J.P. see Kurz, W.A., 909
Kinloch, B.B., Jr., and Dupper, G.E. Restricted distribution of a virulent race of the white pine blister rust pathogen in the western United States, 448
Kinn, D.N. Incidence of pinewood nematode dauerlarvae and phoretic mites associated with long-horned beetles in central Louisiana, 187
Klein, R.M. see Perkins, T.D., 565
Klinka, K. see Kabzems, R.D., 1557
Klinka, K. see Kabzems, R.D., 1565
Knowles, P. see Barrett, J.W., 379
Knowles, P.H. see Barrett, J.W., 756
Kozlowski, T.T. see Yamamoto, F., 69
Kriebel, H.B. see Whitmore, F.W., 408
Kropp, B. see Goodell, B., 1552
Kukkonen, J. see Raunemaa, T., 466
Kulkarni, D.K. see Khuspe, S.S., 1361
Kurz, W.A., and Kimmins, J.P. Analysis of some sources of error in methods used to determine fine root production in forest ecosystems: a simulation approach, 909
Lalonde, M. see Bousquet, J., 219
Lambeth, C.C. see Foster, G.S., 157
Langlois, C.G. see Gagnon, J., 840
Lanner, R.M. see Connor, K.F., 269
Lapp, M.S. see von Rudloff, E., 1013
Larsen, M.J. see Harvey, A.E., 58
Larsen, M.J. see Jurgensen, M.F., 1283
Last, F.T. see Wilson, J., 957
Leak, W.B. Fifty years of compositional change in deciduous and coniferous forest types in New Hampshire, 388
Leak, W.B. Comparison of standard and actual tree-growth trends for deciduous and coniferous species in New Hampshire, 1297
Leary, R. see Hamlin, D., 353
Leduc, D.J. A comparative analysis of the reduced major axis technique of fitting lines to bivariate data, 654
Leduc, D.J. Erratum: A comparative analysis of the reduced major axis technique of fitting lines to bivariate data, 1630

Lennox, T.L., and Lumis, G.P. Evaluation of physical properties of several growing media for use in aerial seeding containers, 165
Lester, D.T. see Thomson, T.A., 510
Letouzey, J. see Comps, B., 1043
Lieffers, V.J. Leader size and number of leaf primordia as factors in timing of terminal bud flush in black spruce and tamarack, 1390
Lieffers, V.J., and Rothwell, R.L. Effects of drainage on substrate temperature and phenology of some trees and shrubs in an Alberta peatland, 97
Lienkaemper, G.W., and Swanson, F.J. Dynamics of large woody debris in streams in old-growth Douglas-fir forests, 150
Linder, S., Benson, M.L., Myers, B.J., and Raison, R.J. Canopy dynamics and growth of *Pinus radiata*. I. Effects of irrigation and fertilization during a drought, 1157
Lindström, A., and Nyström, C. Seasonal variation in root hardiness of container-grown Scots pine, Norway spruce, and lodgepole pine seedlings, 878
Livingston, N.J., and Black, T.A. Water stress and survival of three species of conifer seedlings planted on a high elevation south-facing clear-cut, 1115
Livingston, N.J., and Black, T.A. Stomatal characteristics and transpiration of three species of conifer seedlings planted on a high elevation south-facing clear-cut, 1273
Loewenstein, H. see Entry, J.A., 735
Löfgren, K.G. A fundamental inequality for the assessment of forest land values, 1309
Loo-Dinkins, J.A., and Tauer, C.G. Statistical efficiency of six progeny test field designs on three loblolly pine (*Pinus taeda* L.) site types, 1066
Loopstra, C. see Alden, J., 1519
Lowell, K.E., and Mitchell, R.J. Stand growth projection: simultaneous estimation of growth and mortality using a single probabilistic function, 1466
Lumis, G.P. see Lennox, T.L., 165
Lundmark, T., and Hällgren, J.-E. Effects of frost on shaded and exposed spruce and pine seedlings planted in the field, 1197
Luxmoore, R.J. see O'Neill, E.G., 878
Maass, S.A. see Risenhoover, K.L., 357
Macadam, A.M. Effects of broadcast slash burning on fuels and soil chemical properties in the Sub-boreal Spruce Zone of central British Columbia, 1577
MacDougall, R.G., Thompson, R.G., and Piene, H. Stem electrical capacitance and resistance measurements as related to total foliar biomass of balsam fir trees, 1071
Maclauchlan, L.E., Borden, J.H., Cackette, M.R., and D'Auria, J.M. A rapid, multisample technique for detection of trace elements in trees by energy-dispersive X-ray fluorescence spectroscopy, 1124
Magnussen, S., and Yeatman, C.W. Early testing of jack pine. I. Phenotypic response to spacing, 453
Magnussen, S., and Yeatman, C.W. Early testing of jack pine. II. Variance and repeatability of stem and branch characters, 460
Maguire, C.C. see Maguire, D.A., 1627
Maguire, D.A., and Hann, D.W. Equations for predicting sapwood area at crown base in southwestern Oregon Douglas-fir, 236
Maguire, D.A., and Maguire, C.C. Analyzing treatment effects on Douglas-fir seedling survival with an extreme value regression model, 1627
Mahendrappa, M.K. Tree species and urea treatment effects on sulfur and metals in throughfall and stemflow of some eastern Canadian forest stands, 1035
Marshall, J.D., and Perry, D.A. Basal and maintenance respiration of mycorrhizal and nonmycorrhizal root systems of conifers, 872
Marshall, J.D. see Espinosa Bancalari, M.A., 174
Marshall, P.L., and Jahraus, K. Sample size for foliar analyses of coastal Douglas-fir, 1240
Martell, D.L., Otukol, S., and Stocks, B.J. A logistic model for predicting daily people-caused forest fire occurrence in Ontario, 394
Martin, J.A. see Thomson, T.A., 510
Mascarenhas, A.F. see Khuspe, S.S., 1361
Mason, P.A. see Wilson, J., 957
Massicotte, H.B., Ackerley, C.A., and Peterson, R.L. The root–fungus interface as an indicator of symbiont interactions in ectomycorrhizae, 846
Mattson, K.M., Swank, W.T., and Waide, J.B. Decomposition of woody debris in a regenerating, clear-cut forest in the Southern Appalachians, 712
Maxwell, B.D. see Petersen, T.D., 1620
McAfee, B.J., and Fortin, J.A. The influence of pH on the competitive interactions of ectomycorrhizaql mycobionts under field conditions, 859
McCreary, D.D., and Zaerr, J.B. Root respiration has limited value for assessing Douglas-fir seedling quality, 1144
McCullough, D.G., and Wagner, M.R. Evaluation of four techniques to assess vigor of water-stressed ponderosa pine, 138
McCullough, K. see Pang, P.C., 1379
McCurdy, W.D., and Powell, G.R. Syllepsis in *Larix laricina*: association of sylleptic branching with cross-sectional stem growth and stem form of saplings, 1609
McGill, W.B. see Fyles, J.W., 109
McGill, W.B. see Fyles, J.W., 242
McGregor, M.D., Amman, G.D., Schmitz, R.F., and Oakes, R.D. Partial cutting lodgepole pine stands to reduce losses to the mountain pine beetle, 1234
McKee, Jr., W.H. see McKevlin, M.R., 1257
McKee, Jr., W.H. see McKevlin, M.R., 1572

McKevlin, M.R., Hook, D.D., McKee, W.H., Jr., Wallace, S.U., and Woodruff, J.R. Loblolly pine seedling root anatomy and iron accumulation as affected by soil waterlogging, 1257
McKevlin, M.R., Hook, D.D., McKee, W.H., Jr., Wallace, S.U., and Woodruff, J.R. Phosphorus allocation in flooded loblolly pine seedlings in relation to iron uptake, 1572
McLaughlin, R. see Gillingham, M.P., 146
McTague, J.P., and Bailey, R.L. Simultaneous total and merchantable volume equations and a compatible taper function for loblolly pine, 87
Mehta, U. see Khuspe, S.S., 1361
Melachrinoudis, E., Rumpf, D., and Venegas, R. Mixed integer programming improves spray operation planning, 1602
Mendoza, G.A. Goal programming formulations and extensions: an overview and analysis, 575
Meng, C.H., and Tang, S.Z. Interval estimation of operational availability of tree harvesting machines, 1451
Merkle, S.A., and Adams, W.T. Patterns of allozyme variation within and among Douglas-fir breeding zones in southwest Oregon, 402
Merzeau, D. see Comps, B., 1043
Mitchell, R.J., Garrett, H.E., Cox, G.S., Atalay, A., and Dixon, R.K. Boron fertilization, ectomycorrhizal colonization, and growth of *Pinus echinata* seedlings, 1153
Mitchell, R.J. see Baser, C.M., 36
Mitchell, R.J. see Lowell, K.E., 1466
Mittal, R.K., and Wang, B.S.P. Fungi associated with seeds of eastern white pine and white spruce during cone processing and seed extraction, 1026
Molina, R. see Perry, D.A., 929
Moore, T.R. The effect of simulated acid rain on the nutrient status of subarctic woodland soils of eastern Canada, 370
Morgenstern, E.K. see Villeneuve, M., 1147
Morgenstern, E.K. see Villeneuve, M., 1219
Morgenstern, E.K. see Wanyancha, J.M., 1246
Morgenstern, E.K. see Wanyancha, J.M., 1251
Mothe, F. see Corriveau, A., 675
Mroz, G.D. see Becker, C.A., 813
Munro, R.C. see Wilson, J., 957
Myers, B.J. see Linder, S., 1157
Nagmani, R. see von Aderkas, P., 1293
Nambiar, E.K.S. Do nutrients retranslocate from fine roots?, 913
Nautiyal, J.C., and Rawat, J.K. Tenure effects on forest management investment, 284
Navratil, S. see Bella, I.E., 1289
Nebeker, T.E. see Paine, T.D., 1202
Nemani, R.R. see Running, S.W., 472
Neumann, R.W. see Fisher, J.T., 47
Newton, M. see Cole, E.C., 181
Nilsson, J.-E., and Andersson, B. Performance in freezing tests and field experiments of full-sib families of *Pinus sylvestris* (L.), 1340
Norby, R.J. see O'Neill, E.G., 878
Northway, S. see Gillingham, M.P., 146
Nusser, S.M. see Byrne, S.V., 23
Nyland, R. D. see Hansen, G. D., 1
Nyström, C. see Lindström, A., 787
Oakes, R.D. see McGregor, M.D., 1234
Oehlschlager, A.C. see Borden, J.H., 118
Ohsato, S., and Okawara, S. Tensile force required to pull the main line during out haul in lateral extraction using the Gandai cable logging system, 524
Okawara, S. see Ohsato, S., 524
Omi, S.K. see Duryea, M.L., 1371
O'Neill, E.G., Luxmoore, R.J., and Norby, R.J. Increases in mycorrhizal colonization and seedling growth in *Pinus echinata* and *Quercus alba* in an enriched CO_2 atmosphere, 878
O'Reilly, C., and Owens, J.N. Long-shoot bud development, shoot growth, and foliage production in provenances of lodgepole pine, 1421
Otukol, S. see Martell, D.L., 394
Owens, J.N. see O'Reilly, C., 1421
Paine, T.D., and Stephen, F.M. Influence of tree stress and site quality on the induced defense system of loblolly pine, 569
Paine, T.D., Blanche, C.A., Nebeker, T.E., and Stephen, F.M. Composition of loblolly pine resin defenses: comparison of monoterpenes from induced lesion and sapwood resin, 1202
Pang, P.C., Barclay, H.J., and McCullough, K. Aboveground nutrient distribution within trees and stands in thinned and fertilized Douglas-fir, 1379
Park, Y.S., and Fowler, D.P. Genetic variances among clonally propagated population of tamarack and the implications for clonal forestry, 1175
Parker, V.T. see Yoder-Williams, M.P., 991
Parker, W.H. see Van Damme, L., 421
Parresol, B.R., Hotvedt, J.E., and Cao, Q.V. A volume and taper prediction system for bald cypress, 250

Pastor, J., Gardner, R.H., Dale, V.H., and Post, W.M. Successional changes in nitrogen availability as a potential factor contributing to spruce declines in boreal North America, 1394
Patton, R.F. see Stanosz, G.R., 995
Patton, R.F. see Stanosz, G.R., 1001
Payandeh, B., and Ek, A.R. Erratum: Distance methods and density estimators, 95
Perkins, T.D., Vogelmann, H.W., and Klein, R.M. Changes in light intensity and soil temperature as a result of forest decline on Camels Hump, Vermont, 565
Perry, D.A., Molina, R., and Amaranthus, M.P. Mycorrhizae, mycorrhizospheres, and reforestation: current knowledge and research needs, 929
Perry, D.A., Choquette, C., and Schroeder, P. Nitrogen dynamics in conifer-dominated forests with and without hardwoods, 1434
Perry, D.A. see Espinosa Bancalari, M.A., 174
Perry, D.A. see Espinosa Bancalari, M.A., 722
Perry, D.A. see Marshall, J.D., 872
Perry, D.A. see Amarantus, M.P., 944
Petersen, T.D., and Maxwell, B.D. Water stress of *Pinus ponderosa* in relation to foliage density of neighboring plants, 1620
Peterson, R.L. see Massicotte, H.B., 846
Pharis, R.P. see Williams, D.J., 1442
Piene, H. see MacDougall, R.G., 1071
Pierce, H.D., Jr. see Borden, J.H., 118
Pierce, H.D., Jr. see Brooks, J.E., 740
Pineau, M. see Gagnon, R.R., 105
Pitcher, D.C. Fire history and age structure in red fir forests of Sequoia National Park, California, 582
Pope, P.E. see Reinsvold, R.J., 964
Pope, P.E. see Simmons, G.L., 970
Post, W.M. see Pastor, J., 1394
Powell, G.R. Syllepsis in *Larix laricina*: analysis of tree leaders with and without sylleptic long shoots, 490
Powell, G.R. see McCurdy, W.D., 1609
Powell, J.M. see Jozsa, L.A., 1075
Price, C. On a social discount rate for forestry: Comment, 1472
Puri, S., and Tak, M.A. *Accelerating growth of silver fir seedlings under controlled environments, 835*
Puttonen, P. Abscisic acid concentration in Douglas-fir needles in relation to lifting date, cold storage, and postplanting vigor of seedlings, 383
Quinby, P.A. An index to fire incidence, 731
Radamaker, G.K. see Higgins, S.S., 1364
Rai, A.K. see Gunn, E.A., 1507
Raison, R.J. see Linder, S., 1157
Rajora, O.P. see Hyun, J.O., 1134
Raunemaa, T., Hari, P., Kukkonen, J., Anttila, P., and Katainen, H.-S. Long-term changes of needle litter in coniferous forests of Finland, 466
Rawat, J.K. see Nautiyal, J.C., 284
Raynal, D.J. see Jones, R.H., 539
Reed, W.J., and Errico, D. Techniques for assessing the effects of pest hazards on long-run timber supply, 1455
Reinsvold, R.J., and Pope, P.E. Combined effect of soil nitrogen and phosphorus on nodulation and growth of *Robinia pseudoacacia*, 964
Rinne, P., Kauppi, A., and Ferm, A. Induction of adventitious buds and sprouts on birch seedlings (*Betula pendula* Roth and *B. pubescens* Ehrh.), 545
Ripalta, A. see Villalba, R., 1527
Risenhoover, K.L., and Maass, S.A. The influence of moose on the composition and structure of Isle Royale forests, 357
Robertson, A. The centroid of tree crowns as an indicator of abiotic processes in a balsam fir wave forest, 746
Robertson, W.J. see Boyle, C.D., 1480
Rohr, R., and Hanus, D. Vegetative propagation of wavy grain sycamore maple, 418
Rothwell, R.L. see Lieffers, V.J., 97
Rumpf, D. see Melachrinoudis, E., 1602
Running, S.W., Nemani, R.R., and Hungerford, R.D. Extrapolation of synoptic meteorological data in mountainous terrain and its use for simulating forest evapotranspiration and photosynthesis, 472
Rustagi, K.P., and Bare, B.B. Resolving multiple goal conflicts with interactive goal programming, 1401
Rutherford, G.K. see Thacker, D.J., 1138
Rutherford, T.A., and Webster, J.M. Distribution of pine wilt disease with respect to temperature in North America, Japan, and Europe, 1050
Ryker, L.C. see Borden, J.H., 118
Sachs, D. see Sollins, P., 1585
Salonius, P.O. see Boyle, C.D., 1480
Santantonio, D., and Grace, J.C. Estimating fine-root production and turnover from biomass and decomposition data: a compartment–flow model, 900
Santantonio, D., and Santantonio, E. Effect of thinning on production and mortality of fine roots in a *Pinus radiata* plantation on a fertile site in New Zealand, 919

Santantonio, E. see Santantonio, D., 919
Schär, E. see Kienast, F., 683
Schier, G.A. Throughfall chemistry in a red maple provenance plantation sprayed with "acid rain," 660
Schier, G.A. Germination and early growth of four pine species on soil treated with simulated acid rain, 1190
Schlentner, R.E. see Gordon, A.M., 304
Schmitz, R.F. see McGregor, M.D., 1234
Schroeder, P. see Perry, D.A., 1434
Schweingruber, F.H. see Kienast, F., 683
Scrivani, J.A. see Hann, D.W., 17
Sebastian, L.P. see Villeneuve, M., 1147
Sebastian, L.P. see Villeneuve, M., 1219
Shaw, C.G., III, Sidle, R.C., and Harris, A.S. Evaluation of planting sites common to a southeast Alaska clear-cut. III. Effects of microsite type and ectomycorrhizal inoculation on growth and survival of Sitka spruce seedlings, 334
Shaw, C.G., III. see Sidle, R.C., 340
Sidle, R.C., and Shaw, C.G., III. Evaluation of planting sites common to a southeast Alaska clear-cut. IV. Nutrient levels in ectomycorrhizal Sitka spruce seedlings, 340
Sidle, R.C. see Shaw, C.G., III, 334
Simmons, G.L., and Pope, P.E. Influence of soil compaction and vesicular–arbuscular mycorrhizae on root growth of yellow poplar and sweet gum seedlings, 970
Simon, J.-P. see Desponts, M., 1006
Skroch, W.A. see Warren, S.L., 1092
Slagle, R.L. see Hobbs, S.D., 40
Smith, B.G. see Harmon, M.E., 1265
Smith, V.G., and Watts, M. An assessment of the structural method of deriving a black spruce site equation, 1181
Smith, V.G., Watts, M., and James, D.F. Mechanical stability of black spruce in the clay belt region of northern Ontario, 1080
Smith, W.K. see DeLucia, E.H., 527
Sollins, P., Cline, S.P., Verhoeven, T., Sachs, D., and Spycher, G. Patterns of log decay in old-growth Douglas-fir forests, 1585
Speyer, M.R. see Gillingham, M.P., 146
Sprinz, P.T., and Burkhart, H.E. Relationships between tree crown, stem, and stand characteristics in unthinned loblolly pine plantations, 534
Spycher, G. *see Sollins, P., 1585*
Stafford, S.G. see Hobbs, S.D., 40
Stafford, S.G. see Entry, J.A., 865
Stanosz, G.R., and Patton, R.F. Armillaria root rot in Wisconsin aspen sucker stands, 995
Stanosz, G.R., and Patton, R.F. Armillaria root rot in aspen stands after repeated short rotations, 1001
Starbuck, C.J. see Baser, C.M., 36
Stark, N.M. see Entry, J.A., 735
Steenackers, V. see Ceulemans, R., 273
Stephen, F.M. see Paine, T.D., 569
Stephen, F.M. see Paine, T.D., 1202
Stocks, B.J. Fire behavior in immature jack pine, 80
Stocks, B.J. see Martell, D.L., 394
Stoddart, J. see Kim, Y.T., 27
Strong, T.F., and Grigal, D.F. Site and seed source influence jack pine yields in the Lake States, 705
Sucoff, E.I. see Andersen, C.P., 951
Sullivan, T.P., and Vyse, A. Impact of red squirrel feeding damage on spaced stands of lodgepole pine in the Cariboo Region of British Columbia, 666
Sutton, R.F. Root growth capacity and field performance of jack pine and black spruce in boreal stand establishment in Ontario, 794
Swank, W.T. see Mattson, K.M., 712
Swanson, F.J. see Lienkaemper, G.W., 150
Tak, M.A. see Puri, S., 835
Tang, S.Z. see Meng, C.H., 1451
Tappeiner, J.C., II, Hughes, T.F., and Tesch, S.D. Bud production of Douglas-fir (*Pseudotsuga menziesii*) seedlings: response to shrub and hardwood competition, 1300
Tauer, C.G. see Loo-Dinkins, J.A., 1066
Tenow, O. see Gref, R., 346
Tesch, S.D. see Tappeiner, J.C., II, 1300
Thacker, D.J., Rutherford, G.K., and VanLoon, G.W. The effects of simulated acid precipitation in the surface horizons of two eastern Canadian forest Podzol soils, 1138
Thompson, R.G. see MacDougall, R.G., 1071
Thomson, A.J. Comparison of lodgepole pine yield tables, 1110
Thomson, T.A., Lester, D.T., and Martin, J.A. Marginal analysis and cost effectiveness in seed orchard management, 510
Thrower, J.S. Growth intercepts for estimating site quality of young white spruce plantations in north central Ontario, 1385
Timmer, V.R., and Armstrong, G. Growth and nutrition of containerized *Pinus resinosa* at exponentially increasing nutrient additions, 644

Tippett, J.T., and Barclay, J.L. Detection of bark lesions caused by *Phytophthora cinnamomi* in *Eucalyptus marginata* with the plant impedance ratio meter and the Shigometer, 1228
Titus, S.J. see Clyde, M.A., 1223
Umbach, D.M., and Davis, D.D. Influence of humidity and temperature during exposure on SO_2-induced leaf necrosis of Virginia pine and river birch, 1213
Ung, C.-H., et Bertrand, F. Analyse qualitative des équations différentielles d'un modèle de croissance pour l'érablière à sucre, 588
Van Cleve, K. see Gordon, A.M., 304
Van Damme, L., and Parker, W.H. Selection of *Picea mariana* for growing space efficiency, 421
van den Driessche, R. Importance of current photosynthate to new root growth in planted conifer seedlings, 776
Van Deusen, P.C. 3-p sampling and design versus model-based estimates, 115
Van Deusen, P.C. Combining taper functions and critical height sampling for unbiased stand volume estimation, 1416
Van Deusen, P.C. Testing for stand dynamics effects on red spruce growth trends, 1487
VanLoon, G.W. see Thacker, D.J., 1138
Veilleux, J.-M. see Doucet, R., 93
Venegas, R. see Melachrinoudis, E., 1602
Verbyla, D. Erratum: Potential predictions bias in regression and discriminant analysis, 352
Verbyla, D.L. Classification trees: a new discrimination tool, 1150
Verhoeven, T. see Sollins, P., 1585
Villalba, R., Boninsegna, J.A., and Ripalta, A. Climate, site conditions, and tree growth in subtropical northwestern Argentina, 1527
Villeneuve, M., Morgenstern, E.K., and Sebastian, L.P. Estimation of wood density in family tests of jack pine and black spruce using the Pilodyn tester, 1147
Villeneuve, M., Morgenstern, E.K., and Sebastian, L.P. Variation patterns and age relationships of wood density in families of jack pine and black spruce, 1219
Vogelmann, H.W. see Perkins, T.D., 565
Vogt, D.J. see Gower, K.T., 630
Vogt, K.A. see Gower, K.T., 630
von Aderkas, P., Bonga, J.M., and Nagmani, R. Promotion of embryogenesis in cultured megagametophytes of *Larix decidua*, 1293
von Rudloff, E., and Lapp, M.S. Chemosystematic studies in the genus *Pinus*. VI. General survey of the leaf oil terpene composition of lodgepole pine, 1013
Vyse, A. see Sullivan, T.P., 666
Wagner, M.R. see McCullough, D.G., 138
Waide, J.B. see Mattson, K.M., 712
Wallace, S.U. see McKevlin, M.R., 1257
Wallace, S.U. see McKevlin, M.R., 1572
Walters, D.K. see Hann, D.W., 17
Wang, B.S.P. see Mittal, R.K., 1026
Wang, C.J.K. see Wilcox, H.E., 884
Wang, C.J.K. see Wilcox, H.E., 976
Wanyancha, J.M., and Morgenstern, E.K. Genetic variation in response to nitrogen fertilizer levels in tamarack families, 1246
Wanyancha, J.M., and Morgenstern, E.K. Genetic variation in response to soil types and phosphorus fertilizer levels in tamarack families, 1251
Warren, S.L., Skroch, W.A., and Hinesley, L.E. Effect of root competition and nitrogen on growth and mineral content of *Abies fraseri*, 1092
Watts, M. see Smith, V.G., 1080
Watts, M. see Smith, V.G., 1181
Webber, J.E., and Yeh, F.C.H. Test of the first-on, first-in pollination hypothesis in coastal Douglas-fir, 63
Weber, M.G. Decomposition, litter fall, and forest floor nutrient dynamics in relation to fire in eastern Ontario jack pine ecosystems, 1496
Webster, J.M. see Rutherford, T.A., 1050
Weetman, G.F. see Keane, M.G., 205
Wentworth, T.R. see Byrne, S.V., 23
Westman, W.E. Aboveground biomass, surface area, and production relations of red fir (*Abies magnifica*) and white fir (*A. concolor*), 311
Whitmore, F.W., and Kriebel, H.B. Expression of a gene in *Pinus strobus* ovules associated with fertilization and early embryo development, 408
Wilcox, H.E., and Wang, C.J.K. Mycorrhizal and pathological associations of dematiaceous fungi in roots of 7-month-old tree seedlings, 884
Wilcox, H.E., and Wang, C.J.K. Ectomycorrhizal and ectendomycorrhizal associations of *Phialophora finlandia* with *Pinus resinosa, Picea rubens,* and *Betula alleghaniensis*, 976
Williams, D.J., Dancik, B.P., and Pharis, R.P. Early progeny testing and evaluation of controlled crosses of black spruce, 1442
Williams, F. Propagation of mature western white pine (*Pinus monticola* Dougl.) by cuttings, 349
Wilson, J., Mason, P.A., Last, F.T., Ingleby, K., and Munro, R.C. Ectomycorrhiza formation and growth of Sitka spruce seedlings on first-rotation forest sites in northern Britain, 957
Wingfield, M.J. see Bedker, P.J., 51
Wolter, K.E. see Yamamoto, F., 69
Woodard, P.M. see Eberhart, K.E., 1207
Woodruff, J.R. see McKevlin, M.R., 1257

Woodruff, J.R. see McKevlin, M.R., 1572
Yamamoto, F., Kozlowski, T.T., and Wolter, K.E. Effect of flooding on growth, stem anatomy, and ethylene production of *Pinus halepensis* seedlings, 69
Yang, K.C., and Hazenberg, G. Geographical variation in wood properties of *Larix laricina* juvenile wood in northern Ontario, 648
Yeatman, C.W. see Magnussen, S., 453
Yeatman, C.W. see Magnussen, S., 460
Yeh, F.C.H. see Webber, J.E., 63
Ying, C.C., and Hunt, R.S. Stability of resistance among *Pinus contorta* provenances to *Lophodermella concolor* needle case, 1596
Ying, C.C. see Hunt, R.S., 594
Yoder-Williams, M.P., and Parker, V.T. Allelopathic interference in the seedbed of *Pinus jeffreyi* in the Sierra Nevada, California, 991
Zaerr, J.B. see McCreary, D.D., 1144
Zala, C. see Gilbert, D.E., 1060
Zsuffa, L. see Hyun, J.O., 1134